

Composition : Tramadol Hydrochloride 50mg Capsule.

Indications : It is indicated for the treatment of moderate to severe pain.

Dosage and administration : Adult & children over 12 years : For acute pain : an initial dose of 100 mg is usually necessary. This can be followed by doses of 50 mg or 100 mg not more frequently than 4 hourly and duration of therapy should be matched to clinical need. The total daily dosage by mouth should not exceed 400 mg. For chronic pain : an initial dose of 50mg is recommended and then subsequent doses are maintained according to pain severity. Or, as directed by the registered physicians.

Side effects : Most commonly side effects are-dizziness, nausea, constipation, headache, drowsiness, vomiting, pruritus, urticaria, tachycardia, dry mouth.

Contraindications : It is contraindicated in patients with known hypersensitivity to tramadol or any other components of this drug. It is also contraindicated in acute intoxication with alcohol, hypnotics, centrally acting analgesics, opioids or psychotropic drugs.

Use in pregnancy and lactation : There are no adequate and well-controlled studies in pregnant women. Tramadol should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Tramadol has been detected in human breast milk. A decision should be made

Kingdol
Capsule



whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Drug interactions : Co-administration of monoamine oxidase inhibitors with tramadol is not recommended. Co-administration of tramadol with carbamazepine results in markedly decreased serum concentrations of tramadol which may reduce analgesic effect.

Precautions : Tramadol should be used with caution in patients with severe renal or hepatic impairment, head injury, increased intracranial pressure or patients in shock or at risk of convulsions. Care should be taken when administering tramadol to patients with existing respiratory depression or to patients taking concomitant CNS depressant drugs.

Packing : Kingdol 3 x 10's capsules in blister pack.