

Albasine Tablet

Composition : Almitrine Bismesylate + Raubasine; (30mg+10mg) Tablet.

Indications : This tablet is indicated for reduction of neurological damage & accelerates recovery after cerebral stroke, Minor age related neurological disorders, some visual disorders related to the circulation, some disorders of the inner ear related to the circulation (hearing loss, dizziness, buzzing sounds in the ear).

Dosage and administration : 1 tablet once or twice daily (taken separately, several hours apart), without exceeding 2 tablets per day. Or, as directed by the registered physicians.

Contraindication : This tablet is contraindicated in patients with Hypersensitivity to any of the ingredients of it or severe hepatic impairment.

Side effects : Weight loss, nausea, sensations of heaviness or burning in the stomach, gastrointestinal disorders, diarrhoea or constipation, insomnia, drowsiness, agitation, anxiety, dizziness, palpitations.

Precautions : Due to the presence of lactose, this drug should not be used in the case of galactosemia, of glucose and galactose malabsorption syndrome or of lactose deficiency (rare metabolic diseases). Do not exceed the recommended dose.

Use in pregnancy and lactation : It is not recommended in pregnancy. It should be used with caution only when the expected benefit to the mother is greater than the possible risk to the

fetas.

Drug interactions : Do not associated with MAO-Is (monoamine oxidase inhibitor), Not to be used concurrently with other almitrine-containing preparations.

Packing : Albasine : 2x14's tablets in alu alu blister pack.