

**Composition :** Tolfenamic Acid 200mg Tablet.

**Indications :** It is indicated for the treatment of acute migraine.

**Dosage and administration : Adults :** 200 mg when the first symptoms of migraine appear. The treatment can be repeated once after 1-2 hours if a satisfactory response is not obtained.

**Children :** A paediatric doses regimen has not yet been established.

Or, as directed by the registered physicians.

**Side effects :** Tolfenamic acid is well tolerated at the recommended dosage. Most commonly side effects are diarrhea, nausea, abdominal pain, vomiting, flatulence, constipation, dyspepsia, pruritus, urticaria, edema, hypertension, cardiac failure, hepatitis, jaundice, headache, tremor, vertigo, dizziness, fatigue.

**Contraindications :** It is contraindicated in patients with known hypersensitivity to Tolfenamic Acid or any other components of this drug. It is also contraindicated in patients with active or history of recurrent peptic ulcer, who have previously shown hypersensitivity reactions (e.g. asthma, rhinitis, urticaria) in response to aspirin or other NSAIDs.

**Use in pregnancy and lactation :** There are no adequate and well-controlled studies in pregnant women. During the first and second trimester of pregnancy, Tolfenamic acid should not be given unless clearly necessary. Like other NSAIDs, Tolfenamic acid should

## Anilic Tablet

not be given in the last trimester. Tolfenamic acid can appear in breast milk in very low concentrations. It should be avoided when breast feeding.

**Drug interactions :** Co-administration of following drugs with tolfenamic acid is not recommended-anticoagulants, diuretics, antihypertensive, lithium, methotrexate, ciclosporin, corticosteroids.

**Precautions :** Tolfenamic acid should be used with caution in patients with history of asthma, gastrointestinal bleeding, ulceration, liver or kidney impairment.

**Packing :** Anilic 3 x 10's tablets in blister pack.