

## Aslor Tablet

**Composition :** Desloratadine 5mg Tablet.

**Indications :** Allergic Rhinitis : Desloratadine is indicated for the relief of the nasal and non-nasal symptoms of allergic rhinitis (seasonal and perennial) in patients 12 years of age or older. Chronic Idiopathic Urticaria : Desloratadine is indicated for the symptomatic relief of pruritus, reduction in the number of hives and size of hives in patients with Chronic Idiopathic Urticaria 12 years of age and older.

**Dosage & administration :** Adults and 12 years of age or older children 5mg once daily. Child 6-11 years: 2.5mg once daily, 1-5 years: 1.25mg once daily.

Patients with liver or renal impairment : A starting dose of one Aslor tablet (Desloratadine 5mg) every other day is recommended. Or, as directed by the registered physician.

**Contraindication :** Desloratadine is contraindicated in patients who are hypersensitive to this medication or to any of its ingredients or to Loratadine.

**Precaution :** The safety and effectiveness of Desloratadine in pediatric patients under 12 years of age has not been established.

**Side effects :** Desloratadine is generally well tolerated. However, dry mouth, fatigue, somnolence and myalgia are commonly reported side effects.

**Use in pregnancy and lactation :** Desloratadine has been assigned to pregnancy category C by the FDA.

**Drug interaction :** No clinically

important drug interactions have been reported.

**Packing :** Aslor tablet : 10 x 10's tablets in blister pack.