

Composition : Amlodipine+Valsartan; (5mg+80mg), (5mg+160mg) & (10mg+160mg) Tablet.

Indications : This tablet is indicated for the treatment of hypertension :

In patients not adequately controlled on monotherapy.

As initial therapy in patients likely to need multiple drugs to achieve their blood pressure goals.

Dosage and administration : Amlodipine is an effective treatment of hypertension in once daily doses of 2.5mg-10mg while valsartan is effective in doses of 80mg-320mg. In clinical trials with amlodipine and valsartan, the antihypertensive effects increased with increasing doses. Or, as directed by the registered physician.

Contraindication : This tablet is contraindicated in patients who are hypersensitive to any component of this product.

Side effects : The most commonly reported adverse effects are lymphadenopathy, palpitations, tachycardia, ear pain, diarrhea, nausea, constipation, dyspepsia, abdominal discomfort, dry mouth, asthenia, headache, insomnia, pruritus etc.

Precaution : Caution should be exercised when administering this drug to patients with severe and mild-to-moderate hepatic impairment, including patients with biliary obstructive disorder.

Use in Pregnancy and lactation :

Pregnancy Category- Amlodipine-C Valsartan-D by FDA.

Drug interaction : Studies with

AV

Film Coated Tablet

Amlodipine : In clinical trials, amlodipine has been safely administered with thiazide diuretics, beta-blockers, angiotensin-converting enzyme inhibitors, long-acting nitrates, sublingual nitroglycerin, digoxin, warfarin, non-steroidal anti-inflammatory drugs, antibiotic and other hypoglycemic drugs. Studies with Valsartan: No clinically significant pharmacokinetic interactions were observed when Valsartan was co-administered with amlodipine, atenolol, cimetidine, digoxin, furosemide, glyburide, hydrochlorothiazide or indomethacin.

Packing : AV-5/80 : 3x10's tablets in blister pack.

AV-10 : 2x14's tablets in blister pack.

AV-5 : 2x14's tablets in blister pack.