

Composition : Ciprofloxacin+ Hydrocortisone Acetate (3mg+10mg)/ml.

Indications: Ciprocort otic suspension is indicated for the treatment of acute otitis externa in adult and paediatric patients due to susceptible strains of pseudomonas aeruginosa, staphylococcus aureus and proteus mirabilis.

Dosage and administration: Adults and children (one year and older) : After cleansing 2-4 drops instilled thrice to four times a day and at night. Or as directed by the physician.

Pregnancy and lactation: No adequate and well controlled studies have been performed in pregnant women. Caution should be exercised when it is used by a pregnant women. It is better not to administer Ciprocort otic suspension during lactation.

Paediatric use: Safety and efficacy of Ciprocort otic suspension have been established in paediatric patients 2 years and older in adequate and well controlled clinical trials.

Precautions: As with other antibiotic preparations, use of this product may result in overgrowth of non- susceptible organisms including fungi. If the infection is not improved after one week of therapy, culture should be obtained to guide future treatment.

Contraindication: It should be discontinued at the first appearance of a skin rash or any other sign of hypersensitivity.

Ciprocort

Sterile Otic Suspension

Side effects: Rare reports of headache, pruritus, migraine, hyperesthesia, paresthesia, fungal dermatitis, cough, rash, urticaria, and alopecia have been reported.

Packing: 10ml in plastic dropper bottle.