

**Composition :** Levocetirizine Dihydrochloride 5mg Tablet.

**Indications :** Clarigen is indicated for the treatment of Seasonal allergic rhinitis, Perennial allergic rhinitis, chronic idiopathic urticaria, Food allergy, Allergic bronchial asthma, Eczema.

**Dosage and administration :** Levocetirizine is indicated for all ages above six years. It should be taken as a single dose with or without food. Adult : One Levocetirizine 5mg tablet once daily. Children > 6 years : One Levocetirizine tablet (5mg) once daily. Dose adjustment is indicated in moderate or severe renal impairment. A reduction in the dose of Levocetirizine by 50% should be considered in patients with moderate to severe renal dysfunction. Or, as directed by the registered physician.

**Contraindication :** Levocetirizine is contraindicated in patients who are hypersensitive to Levocetirizine.

**Side effects :** Levocetirizine is generally well tolerated. However, dry mouth, somnolence, fatigue & myalgia are reported occasionally. Less common side effects may include dizziness, headache, nausea etc. Rarely rash, pruritus & urticaria may occur.

**Use in pregnancy and lactation :** Levocetirizine has been assigned to Pregnancy Category B by the FDA.

**Drug interaction :** No clinically

## Clarigen Tablet

important drug interactions have been reported.

**Packing :** Clarigen Tablet : 10x10's tablets in blister pack.