

Composition : Cefpodoxime + Clavulanic Acid; (100mg+62.5mg) & (200mg+125mg) Tablet.

Indications : It is indicated in the following infections when caused by susceptible organisms: Acute bacterial exacerbations of chronic bronchitis, Acute community acquired pneumonia, Upper and lower respiratory tract infections, Skin and soft tissue infections, Urinary tract infections, Pharyngitis and/or tonsillitis, General gonorrhea (men and women) and rectal gonococcal infections (women), Acute maxillary sinusitis.

Dosage and administration : Cefpodoxime Dosage Schedule :

Type of Infection	Total Daily Dosage	Dosage Frequency	Duration
Pharyngitis and /or tonsillitis	200mg	100mg 12 hrly	5 to 10 days
Acute community acquired pneumonia	400mg	200mg 12 hrly	14 days
Acute bacterial exacerbations of chronic bronchitis	400mg	200mg 12 hrly	10 days
Uncomplicated gonorrhea (men and women) and rectal gonococcal infections (women)	200mg	200mg 24 hrly	Single dose
Skin and skin structure infections	800mg	400mg 12 hrly	7 to 14 days
Acute maxillary sinusitis	400mg	200mg 12 hrly	10 days
Uncomplicated urinary tract infection	200mg	100mg 12 hrly	7 days

Or, as directed by the registered physician.

Contraindication : It is contraindicated in patients with known hypersensitivity to the drug.

Side effects : Side effects are diarrhea, nausea, abdominal pain, headache, painful or urination, unusual weakness, easy bleeding and bruising, confusion, dry mouth, chills, body aches, fever, sore throat.

Precautions : Cefpodoxime, like other cephalosporins, should be administered with caution to patients receiving concurrent treatment with potent diuretics. In patients with renal impairment, dosage should be adjusted.

Clavuran Tablet

Use in pregnancy and lactation : Cefpodoxime has been assigned to pregnancy category B by the FDA.

Drug Interactions : Concomitant administration of high doses of antacids (Sodium Bicarbonate and Aluminium Hydroxide) or H₂ blockers reduces peak plasma level by 24% to 42% and the extent of absorption by 27% to 32% respectively.

Packing :

Clavuran-100 : 2x7's tablets in blister pack.

Clavuran-200 : 2x7's tablets in blister pack.