**Composition: Tamsulosin** 

Hydrochloride modified release &
Dutasteride immediate release pellets
ph. Gr. eqv to Tamsulosin
Hydrochloride USP O.4mg &
Dutasteride INN O.5mg.

Indications: This Capsule is indicated for the treatment of the signs and symptoms of benign prostatic hyperplasia (BPH) in men with an enlarged prostate.

Dosage and administration: The recommended dose of this capsule is one capsule once daily approximately 30 minutes after the same meal each day. This capsule should be swallowed whole and should not be

crushed or chewed. No dosage adjustment is required for subjects with renal impairment or for the elderly. Or, as directed by the registered physicians.

Contraindication: This Capsule is contraindicated for use by women and children. It is contraindicated in patients known to be hypersensitive to tamsulosin HCl, dutasteride or any component of the capsule.

Use in Pregnancy and lactation: This Capsule is not indicated for use in woman. These Capsules are not indicated for use in nursing mothers.

Side effects: The following adverse (1. reactions have been reported during the use of this capsule dizziness, abnormal ejaculation, back pain, chest pain, diarrhea, tooth disorder, libido decreased, insomnia, rhinitis, pharyngitis, sinusitis, and asthenia.

## **Combomax**

Capsule

Precautions: lower urinary tract symptoms of BPH can be indicative of other urological diseases, including prostate cancer. Patients should be assessed to rule out other urological 'diseases prior to treatment with this capsule. The treatment of severely renal impaired patients (creatinine clearance is less than 10mll min) should be approached with caution as these patients have not been studied.

Drug interactions: Concurrent administration of other alpha 1 adrenoceptor antagonists could lead to hypotensive effects. This capsule should not be used in combination yvith other alpha-adrenergic blocking agents.

other alpha-adrenergic blocking agents. This tablet should be used with caution in combination with cimetidine. Caution should be exercised with concomitant administration of warfarin and this capsule,

Packing: Combomax: Each box contains 4 x 7's capsules in blister pack.