

Daparol Tablet

Composition : Dapagliflozin 5 mg Tablet.

Indications : Dapagliflozin is indicated as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus .

Dosage & Administration : The recommended starting dose is 5 mg once daily, taken in the morning, with or without food. It may increase to 10 mg orally once a day in patients tolerating therapy with 5 mg and requiring additional glycemic control. Or, as directed by the registered physicians.

Side effects : The most common reported side effects are dehydration, hypotension, impairment in renal function, hypoglycemia, genital mycotic infections, increase in bad cholesterol (LDL-C) and bladder cancer .

Contraindications : Dapagliflozin is contraindicated in patients with hypersensitivity to Dapagliflozin or any component of this medications.

Precaution : Dapagliflozin is not recommended for patients with type 1 diabetes mellitus or for the treatment of diabetic ketoacidosis. It should not be used in patient with active bladder cancer and who having severe kidney disease.

Pregnancy and lactation : Pregnancy Category-C. There are no adequate and well-controlled studies in pregnant women. Dapagliflozin should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Lactation- It is not known whether this drug is excreted in human milk. Caution should be exercised when

Dapagliflozin tablet is administered to a nursing woman.

Drug Interactions : May Enhance the adverse or toxic effect of other hypoglycemic agents, hypotensive agents, Loop Diuretics, MAO inhibitors and Selective Serotonin Reuptake Inhibitors with dapagliflozin.

Packing : Daparol : 1 x 14's tablets in blister pack.