

Composition : Amoxicillin 250mg, 500mg Capsule, 125mg/5ml Suspension, 250mg / 5ml DS Suspension.

Indications : Urinary tract infections : nephritis, pyelonephritis, pyelitis, cystitis, gonorrhoea and non-gonococcal urethritis.

Skin and soft tissue infections : cellulitis, wounds, carbuncles and furunculosis.

Gastro-intestinal infections : enteric fever, ear, nose and throat infections, meningitis.

Dosage & administration : Oral : Standard adult dosage : 250mg three times daily, increasing to 500mg three times daily for more severe infections.

Children (2 to 10 years) : 125mg three times daily, increasing to 250mg three times daily for more severe infection.

Infants (Up to 2 years) : 62.5 - 125mg three times daily or as directed by the physician. In case of severity the dosage may safely be increased. Food does not interfere with absorption of Demoxil.

Side effects : No untoward side-effects have been reported except in some cases which are rare, mild and transitory. Most common side-effects are different types of skin rashes & diarrhoea.

Contraindication : The use of Demoxil is contra-indicated in patients with a history of allergic reaction to penicillin.

Precautions : It should be used with caution in patients with impaired renal function and infections of mononucleosis.

Use in pregnancy and lactation :

Demoxil

Cap. / Suspension

Amoxicillin has been assigned to pregnancy category B by the FDA.

Packing :

Demoxil 250mg Capsule : 5x10 capsules in blister pack.

Demoxil 500mg Capsule : 5x10 capsules in blister pack.

Demoxil : 100ml suspension.

Demoxil-DS : 100ml suspension.