

Pancil Tablet

Composition : Pancil Tablet: Each Tablet Contains Cilostazol USP 100mg.

Indications : Cilostazol is indicated for the reduction of symptoms of intermittent claudication, as indicated by an increased walking distance.

Dosage and administration : The recommended dosage of Cilostazol is 100 mg twice daily taken at least half an hour before or two hours after breakfast and dinner. A dose of 50 mg twice daily should be considered during co-administration of ketoconazole, Itraconazole, Fluconazole, Omeprazole, Erythromycin, Diltiazem. Or as directed by the registered physicians.

Side effects : The most common side effects are headache, dizziness, diarrhea, nausea, vomiting, flatulence, abdominal pain, rhinitis, rash.

Contraindication : Cilostazol is contraindicated in patients with congestive heart failure of any severity. Cilostazol is also contraindicated in patients with hypersensitivity to Cilostazol or any other component of this medications.

Use in pregnancy and lactation : Pregnancy Category-C. There are no adequate and well-controlled studies in pregnant women. It is excreted in human milk. Because of the potential risk to nursing infants, a decision should be made to discontinue nursing or to discontinue Cilostazol.

Drug interactions : Pharmacokinetic studies have shown that Erythromycin, Omeprazole increase the systemic exposure of Cilostazol and/or its major metabolites.

Precautions : Cilostazol should be used with special caution in patients with severe renal impairment and hepatic impairment.

Packing : Pancil Tablet : Each box contains 2 x 14's tablets in blister pack.