

Ejalong Tablet

Composition : Dapoxetine 30mg & 60mg Tablet.

Indications : It is indicated for the treatment of premature ejaculation (PE) in adult men aged 18 to 64 years of age.

Dosage and administration : Adult men (aged 18 to 64 years): The recommended starting dose for all patients is 30 mg, taken as needed approximately 1 to 3 hours prior to sexual activity. If the individual response to 30 mg is insufficient and the patient has not experienced moderate or severe adverse reactions, the dose may be increased to a maximum recommended dose of 60 mg taken as needed approximately 1 to 3 hours prior to sexual activity. Or, as directed by the registered physician.

Side effects : The most common side effects of this tablet are- nausea, dizziness, headache, diarrhoea, insomnia, fatigue, anxiety, restlessness, somnolence, disturbance in attention, tremor, blurred vision, tinnitus, vomiting, constipation, abdominal pain, dyspepsia, flatulence, stomach discomfort, abdominal distension, dry mouth, erectile dysfunction, libido decreased.

Contraindication : It is contraindicated in patients with a history of hypersensitivity to Dapoxetine or any other components of this product.

Use in pregnancy and lactation : This tablet is not indicated for women.

Drug interactions : Concomitant use of alcohol and dapoxetine increases the chance or severity of adverse reactions such as dizziness, drowsiness, slow reflexes, or altered judgment. Caution is advised when dapoxetine is used in

patients taking warfarin chronically. Concomitant use of dapoxetine and potent CYP3A4 inhibitors, such as ketoconazole, itraconazole, ritonavir, saquinavir, telithromycin, nefazodone, nelfinavir and atazanavir, may reduce dapoxetine clearance. Concomitant treatment with moderate CYP3A4 inhibitors (e.g., erythromycin, clarithromycin, fluconazole, amprenavir, fosamprenavir, aprepitant, verapamil, diltiazem) may give rise to significantly increased exposure of dapoxetine.

Precautions : This tablet should not be prescribed to men who have not been diagnosed with Premature Ejaculation. It should not be used in men with erectile dysfunction (ED) who are using PDE5 inhibitors. It is not recommended for use in patients with severe renal impairment. This product should be used with caution in patients with raised intraocular pressure or those at risk of angle closure glaucoma. Possibly prodromal symptoms such as nausea, dizziness/lightheadedness and diaphoresis were reported more frequently among patients treated with this drug. Caution is advised in patients taking this tablet, particularly in concomitant use with medicinal products known to affect platelet function as well as in patients with a history of bleeding or coagulation disorders.

Packing : Ejalong 30 : 1 x 6's tablets in blister pack.

Ejalong 60 : 1 x 6's tablets in blister pack.