

Composition : Fexofenadine Hydrochloride 60mg, 120mg, 180mg Tablet & 30mg/5ml Suspension.

Indications : Fexofenadine Hydrochloride is indicated for the relief from physical symptoms associated with seasonal allergic rhinitis and treatment of chronic idiopathic urticaria. It is not cure totally but prevents the aggravation of rhinitis and urticaria. It reduces the severity of the symptoms and providing much relief from repeated sneezing, runny nose, itchy eyes and general body fatigue caused by rhinitis and urticaria.

Dosage and administration : Fexofast tablets: Seasonal allergic rhinitis and chronic idiopathic urticaria: Adults and Children 12 Years and Older: The recommended dose of Fexofenadine Hydrochloride is 60 mg twice daily or 180 mg once daily with water. A dose of 60 mg once daily is recommended as the starting dose in patients with decreased renal function. Children 6 to 11 Years: 30 mg twice daily with water. A dose of 30 mg once daily is recommended as the starting dose in patients with decreased renal function.

Oral suspension: Seasonal allergic rhinitis: Children 2 to 11 years: The recommended dose of oral suspension 30 mg twice daily. If paediatric patients decrease renal function, then adjust or maintain dose of 30 mg (5 ml) once daily recommended as the starting dose.

Chronic idiopathic urticaria: Children 6 months to 11 years: 30 mg (5 ml) twice daily for patients 2 to 11 years of age and 15 mg (2.5ml) twice daily for patients 6 months to less than 2 years of age. For paediatric patients with decrease renal function the recommended starting dose of oral suspension is 30 mg (5ml) once daily for patients 2 to 11 years of age and 15 mg (2.5ml) once daily for patients 6 months to less than 2 years of age. Shake bottle well, before each use. Or, as directed by the registered physician.

Fexofast

Tablet / Suspension

Use in pregnancy and lactation : Fexofenadine has been assigned to pregnancy category C by the FDA.

Side effects : Nausea, dizziness, vomiting, weakness, drowsiness, sleepiness, fatigue, diarrhea, unusual bowel movements, headache & fever.

Contraindication : Fexofenadine Hydrochloride is contraindicated in patients who are hypersensitive to this medication or its ingredients.

Precautions : Fexofenadine Hydrochloride should generally be avoided in patients with hepatic insufficiency. Patients with renal insufficiency should be given a lower initial dose.

Drug interactions : Fexofenadine Hydrochloride should not be taken closely in time with aluminum and magnesium containing antacids. Co-administration of fexofenadine hydrochloride with either ketoconazole or erythromycin lead to increased plasma concentrations of fexofenadine in healthy adult subjects. Fexofenadine had no effect on the pharmacokinetics of either erythromycin or ketoconazole.

Packing :

Fexofast-60 : 5x 10's tablets in blister pack.

Fexofast-120 : 3x 10's tablets in blister pack.

Fexofast-180 : 2x10's tablets in blister pack.

Fexofast Suspension : 50ml suspension.