

**Composition :** Moxifloxacin 5mg/ml as Moxifloxacin Hydrochloride.

**Indications and uses :** Floxalone eye drop is indicated for the treatment of bacterial conjunctivitis caused by susceptible strains of the following organisms:

**Aerobic Gram-positive microorganisms:** Corynebacterium species, Micrococcus luteus, Staphylococcus aureus, Staphylococcus epidermidis, Staphylococcus haemolyticus, Staphylococcus hominis, Staphylococcus warneri, Streptococcus pneumoniae, Streptococcus viridans group.

**Aerobic Gram-negative microorganisms:** Acinetobacter lwoffii, Haemophilus influenzae, Haemophilus parainfluenzae.

Other microorganisms: Chlamydia trachomatis.

**Dosage and administration :** The recommended dose is one drop in the affected eye every 8 hours (3 times) daily for 7 days. Or, as directed by the registered physician.

**Contraindication :** It is contraindicated in person with a known hypersensitivity to Moxifloxacin, to other quinolones, or to any components in this medication.

**Side effects :** The most frequently reported ocular adverse events were decreased visual acuity, dry eye, ocular discomfort, ocular hyperemia, ocular pain, ocular pruritus, subconjunctival hemorrhage and tearing. These events occurred in approximately 1-6% of patients.

## Floxalone

### Sterile Eye Drops

**Precautions :** As with other antibiotics, prolonged use may result in overgrowth of non-susceptible organisms, including fungi. If super-infection occurs, discontinue use and introduce alternative therapy.

**Pregnancy and lactation :** Pregnancy Category **C** by FDA.

**Drug interaction :** Drug interaction studies have not been conducted with moxifloxacin hydrochloride solution. In vitro studies indicated that moxifloxacin is unlikely to alter the pharmacokinetics of drugs metabolized by the cytochrome P450 enzymes.

**Packing :** 5ml in plastic dropper bottle.