

Composition : Fluticasone propionate 0.005%.

Indications : Fluticasone propionate is a medium potency corticosteroid indicated for the relief of the inflammatory and pruritic manifestations of corticosteroid-responsive dermatoses eczema & psoriasis.

Dosage and administration : Apply a thin film of Flutiderm Ointment to the affected skin areas twice daily. Rub in gently. Or as directed by the physician.

Contraindication : Rosacea, acne and peri-oral dermatitis. Skin lesions caused by infection with viruses (e.g. herpes simplex, chickenpox), fungi (e.g. candidiasis, tinea) or bacteria (e.g. impetigo), hypersensitivity to the preparation.

Precautions : Fluticasone propionate ointment should not be used in the treatment of preexisting skin atrophy and where infection is present at the treatment site.

Drug interactions : No data is available.

Side effects : The following local adverse reactions have been reported; dryness, folliculitis, hypopigmentation, perioral dermatitis, allergic contact dermatitis, secondary infection, skin atrophy, striae. Also, there are reports of the development of pustular psoriasis from chronic plaque psoriasis following reduction or discontinuation of potent topical corticosteroid products.

Flutiderm Ointment

Pregnancy and lactation : Pregnancy Category C.

Packing : Flutiderm ointment : 10gm in a tube.