

Composition : Indapamide+Perindopril Erbumine; (1.25mg+2mg) & (1.25mg+4mg) Tablet.

Indications : It is indicated in the treatment of essential hypertension where blood pressure is not adequately controlled on perindopril alone.

Dosage and administration : 1-2 tablet of **Indapa-plus 2** or One tablet of **Indapa-plus 4** per day preferably taken in the morning before meal. In case of renal insufficiency treatment should be start with an adequate dose.

Contraindication : This fixed combination is contraindicated in following situations. Hypersensitivity to perindopril/ indapamide or to any other angiotensin-converting enzyme (ACE) inhibitor.

Side effects : Side-effects include nausea, vomiting, dyspepsia, diarrhea, constipation, headache, dizziness, sleep disturbance, asthenia, dry mouth, hypotension, persistent dry cough, angioedema, rash.

Precautions : The drug should be used cautiously in the following situations- in renal impairment, renal artery stenosis, heart failure, gout, diabetes, liver cirrhosis, in elderly patient, atherosclerosis, idiopathic or hereditary angioedema.

Use in Pregnancy and lactation : There are no adequate and well-controlled studies in pregnant woman. So in pregnancy **Indapa-plus** is not recommended. Mothers taking **Indapa-plus** should not breast feed.

Drug interactions : Concomitant use of antihypertensive & antipsychotic drug may

Indapa-Plus Tablet

potentiate hypotensive effect and NSAIDs may reduce antihypertensive effect. ARB and Potassium Sparing diuretics may cause hyperkalaemia.

Packing : Indapa-Plus 2 : 2x10's tablets in blister pack.

Indapa-Plus 4 : 1x10's tablets in blister pack.