

Composition : Ivabradine 5mg & 7.5mg Tablet.

Indications : Ivabradine is indicated for symptomatic treatment of chronic stable angina pectoris in coronary artery disease adults with normal sinus rhythm, who has contraindications or intolerance with beta-blockers.

Dosage and administration : The usual recommended starting dose of Ivabradine is 5 mg twice daily after 3-4 weeks of treatment, the dose may be increased to 7.5mg twice daily depending on therapeutic response. It may be adjusted to 2.5mg twice daily as necessary. Or, as directed by the registered physician.

Contraindication : Ivabradine is contraindicated in patients with hypersensitivity to any component of this medication.

Side effects : The most common adverse reactions with Ivabradine are luminous phenomena and bradycardia. These effects are dose dependent and related to the pharmacological effect of the medicinal product.

Precautions : Ivabradine is not recommended in patients with atrial fibrillation or other cardiac arrhythmias, severe hypotension, unstable angina, prolong QT syndrome.

Use in pregnancy and lactation : Ivabradine is contra-indicated during pregnancy and lactation.

Drug interactions : The concomitant use of cardiovascular and non cardiovascular QT prolonging medicinal products with Ivabradine should be avoided since QT prolongation may be

Ivaten Tablet

exacerbated by heart rate reduction.

Overdose : The overdose of Ivabradine may cause severe and prolonged bradycardia.

Packing :

Ivaten-5 : 1x10's tablets in blister pack.

Ivaten-7.5 : 1x10's tablets in blister pack.