

## Lansec Capsule

**Composition** : Lansoprazole 15mg & 30mg Capsule.

**Indications** : Healing and long term management of gastroesophageal reflux disease. Healing and maintenance therapy for patients with duodenal ulcer. Relief of reflux-like symptoms (eg. Heartburn) and/or ulcer-like symptoms (eg. upper epigastric pain) associated with acid-related dyspepsia. Healing of benign gastric ulcer. Lansec is also effective in patients with benign peptic lesions, including reflux oesophagitis, unresponsive to H<sub>2</sub> receptor antagonists. Eradication of H.pylori from the upper gastro-intestinal tract in patients with duodenal ulcer or gastritis when used in combination with appropriate antibiotics.

**Side effects** : Lansec is well tolerated, with adverse effects generally being mild and transient. The most commonly reported adverse effects are headache, dizziness, fatigue and malaise. Gastrointestinal effects include diarrhea, constipation, abdominal pain, dyspepsia, nausea, vomiting, flatulence and dry or sore mouths. Dermatological reactions include skin rashes, urticaria and pruritus.

**Drug interactions** : Caution should be exercised when oral contraceptives and preparations such as phenytoin, theophylline or warfarin are taken concomitantly with the administration of Lansec. Antacids and sucralfate may reduce the bioavailability of lansoprazole.

### Dosage and administration :

Indication	Recommended Dose	Frequency
<b>Duodenal Ulcers</b>		
Short-Term Treatment	15mg	Once daily for 4 weeks
Maintenance of Healed	15mg	Once daily
<b>H. pylori Eradication to Reduce the Risk of Duodenal Ulcer Recurrence*</b>		
<b>Triple Therapy :</b>		
Lansec	30mg	Twice daily (q 12h for 10 or 14 days
Demoxil	1 gram	Twice daily (q 12h) for 10 or 14 days
Clarix	500mg	Twice daily (q 12h) for 10 or 14 days
<b>Dual Therapy :</b>		
Lansec	30mg	Three times daily (q 8th) for 14 days
Demoxil	1 gram	Three times daily (q 8th) for 14 days
<b>Benign Gastric Ulcer</b>		
Short-Term Treatment	30mg	Once daily for up to 8 weeks
<b>NSAID-associated Gastric Ulcer</b>		
Healing	30mg	Once daily for 8 weeks
Risk Reduction	15mg	Once daily for up to 12 weeks

### Gastroesophageal Reflux Disease (GERD)

Short-Term Treatment of Symptomatic GERD 15mg  
Once daily for up to 8 weeks

Short-Term Treatment of Erosive Esophagitis 30mg  
Once daily for up to 8 weeks

or, as directed by the registered physicians.

**Contraindication** : The use of Lansec is contraindicated in patients with a history of hypersensitivity to any of the ingredients of Lansec capsules.

**Precautions** : Symptomatic response to therapy with lansoprazole does not preclude the presence of gastric malignancy. Lansec should be taken before eating.

**Use in pregnancy and lactation** : Lansoprazole has been assigned to pregnancy category B by the FDA.

### Packing :

**Lansec-15** : 6 x 10's capsules in blister pack.

**Lansec-30** : 6 x 10's capsules in blister pack.