

Composition : Levofloxacin 5mg / ml.

Indications : Levoflox eye drop is indicated for the treatment of bacterial conjunctivitis caused by susceptible strains of the following organisms: Aerobic gram-positive, gram-negative micro organisms. In some study it has been found that Levofloxacin is indicated in post-operative ocular infection.

Dosage and administration : Day 1 & 2 : Instill 1-2 drops in the affected eye (s) every 2 hours while awake, up to 8 times per day. Days 3 through 7 : Instill 1-2 drops in the affected eye (s) every 4 hours while awake, up to 4 times per day. Or as directed by the registered physician.

Contraindication : Hypersensitivity to Levofloxacin, to other quinolones or to any component in this medication.

Side effects : The most frequently reported adverse reactions are transient decreased vision, fever, foreign body sensation, headache, ocular pain or discomfort, and photophobia. These events occurred in approximately 1-3% of patients included allergic reactions, occurring in less than 1% of patients included allergic reactions, lid edema, ocular dryness, and ocular itching.

Use in pregnancy and lactation : Pregnancy Category **C** by FDA.

Pediatric use : Safety and efficacy in infants below the age of 1 year have not been established.

Precautions : Prolonged use may result in overgrowth of non-susceptible

Levoflox

Sterile eye drops

organisms, including fungi. If super-infection occurs, discontinue use and institute alternative therapy. Patients wearing contact lenses must not use the drops during the time the lenses are worn. Should not be injected subconjunctivally, nor should it be introduced directly into the anterior chamber of the eye.

Packing : 5ml in plastic dropper bottle.