

Linarol Tablet

Composition : Linagliptin 5 mg Tablet.

Indications : Indicated for adults with diabetes mellitus type II along with diet and exercise to lower blood sugar, may be used as monotherapy or in combination with other common antidiabetic medications including metformin, sulfonylurea, pioglitazone, or insulin.

Dosage and administration : Linagliptin 5 mg once daily. If added to metformin, the dose of metformin should be maintained concomitantly, when used in combination with sulfonylurea, a lower dose of sulfonylurea may be considered to reduce the risk of hypoglycemia. Patients with renal impairment no need to dose adjustment required. Linagliptin can be taken with or without a meal at any time of the day. Or, as directed by the registered physician.

Contraindication : Hypersensitivity to the active substance or to any of the excipients.

Side effects : There may be hypoglycaemia, nasopharyngitis, cough and pancreatitis combination with or add on to metformin and sulfonylurea.

Precautions : Linagliptin should not be used in patients with type I diabetes or for the treatment of diabetic ketoacidosis.

Use in Pregnancy and lactation : Linagliptin has been assigned to pregnancy category B by the FDA.

Drug interactions : Linagliptin is a weak competitive and a weak to moderate mechanism based inhibitor of CYP isozyme CYP3A4, but does not inhibit other CYP isozymes. The risk for clinically meaningful interactions by other medicinal products on linagliptin is low and in clinical studies linagliptin had no clinically relevant effect on the pharmacokinetics of metformin, glyburide, simvastatin, warfarin, digoxin or oral contraceptives.

Packing : Linarol : 2 x 14's tablets in blister pack.