

Composition : Pitavastatin 2mg Tablet.

Indications : Pitavastatin is indicated for - as an adjunctive therapy to diet to reduce elevated total cholesterol (TC) , Low-density lipoprotein cholesterol (LDL-C), Apolipoprotein B (Apo B), Triglycerides (TG), and to increase HDL-C in adult patients with primary hyperlipidemia or mixed dyslipidemia.

Dosage and administration : The dose range for Pitavastatin is 1 to 4 mg orally once daily at any time of the day with or without food. The recommended starting dose is 2 mg and the maximum dose is 4 mg. The administration starting dose and maintenance dose should be individualized to patient characteristics, such as goal of therapy and response. Or, as directed by the registered physician.

Contraindications : Pitavastatin is contraindicated in patients with hypersensitivity to any component of this medications, active liver disease or unexplained persistent elevations of hepatic transaminase levels.

Side effects : Pitavastatin is generally well tolerated. The most common side effects are constipation, headache, myalgia, asthenia, abdominal pain and dizziness.

Pregnancy and lactation : Pregnancy Category-X. Not for use in pregnancy.

Drug interactions : Cyclosporine, erythromycin, niacin, rifampin and fibrate may significantly increased Pitavastatin

Liploss

Film Coated Tablet

exposure. In patients taking these medication, a reduction in Pitavastatin dosage should be considered.

Packing : Liploss : 2x10's tablets in blister pack.