

Composition : Mebeverine HCl 135mg Tablet & 200mg SR Capsule.

Indications : Meverine is indicated for the symptomatic treatment of irritable bowel syndrome and other conditions usually included in this grouping such as: chronic irritable colon, spastic constipation, mucous colitis, spastic colitis. Meverine is also effective in colicky abdominal pain and cramps, persistent, non-specific diarrhoea (with or without alternating constipation) and flatulence.

For the symptomatic treatment of gastrointestinal spasm secondary to organic diseases.

Meverine-SR : Mebeverine Hydrochloride is indicated: For the symptomatic treatment of irritable bowel syndrome (IBS), other conditions include: Chronic irritable colon, Spastic constipation, Mucous Colitis, Colicky abdominal pain, Persistent non-specific diarrhoea.

Dosages : Adults & Children over 10 years, 1 tablet 3 times daily, preferably 20 minutes before meals or as directed by a registered physician.

Meverine-SR : Adults, elderly : 1 Capsule twice daily, preferably 20 minutes before meals. Or, as directed by the registered physician.

Precautions : Caution should be exercised in porphyria or allergic reaction to this or any other medicine of this group.

Use in pregnancy and lactation : No teratogenicity has been shown in animal experiments. However, the usual precautions concerning the administration

Meverine

Tablet/Capsule

of any drug during pregnancy should be exercised. Mebeverine dose not excrete in the breast milk after administering at therapeutic dose.

Drug interaction : Not known.

Contraindication : Hypersensitivity to the drug or any other ingredients.

Side effects : Not known.

Packing : Meverine : 5 x 10's tablets in blister pack.

Meverine SR : 3x10's Capsules in blister pack.