

Composition : Miconazole Nitrate USP 2% .

Indications : It is indicated for the treatment of mycotic infections of the skin & nails, superinfection due to gram-positive bacteria and also topically used in the treatment of tinea pedis (athlete's foot), tinea cruris, tinea corporis, cutaneous candidiasis (moniliasis), tinea versicolor, stomatitis angularis, otitis externa.

Dosage and administration : (For all the ages) : It should be usually applied 1 to 2 times daily for 2 to 4 weeks. For skin infections : The cream should be applied twice daily (morning & evening) to the lesions. Treatment should be prolonged for 10 days after all lesions have disappeared to prevent relapse. For nail infections: The cream should be applied once or twice daily to the lesions. Treatment should be prolonged for 10 days after all lesions have disappeared to prevent relapse. Or, as directed by the registered physicians.

Side effects : Occasionally irritation has been reported.

Contraindication : It may be contraindicated in patients known to have hypersensitivity to the cream or any of its components.

Use in pregnancy and lactation : Only small amounts of Miconazole nitrate are adsorbed following local administration. However, as with other imidazoles, Miconazole nitrate should be used with caution during pregnancy.

Drug Interactions : No drug interactions have been reported.

Precaution : It is for external use only. Contact with the eyes should be avoided at the use of this product. Keep out of reach of children.

Micoderm Cream

Packing : Micoderm Cream : Each laminated tube contains 10gm cream.