

Composition : Miconazole 2%.

Indications : This gel is indicated for oral treatment and prevention of fungal infections of the oropharynx and of super infections due to Grampositive bacteria.

Dosage and administration : For localized lesions of the mouth, a small amount of gel may be applied directly to the affected area with a clean finger. For topical treatment of the oropharynx, the gel should be kept in the mouth for as long as possible. Treatment should be continued for up to 2 days after the symptoms have cleared. For oral candidosis, dental prostheses should be removed at night and brushed with the gel. Or, as directed by the registered physician.

Contraindication : Known hypersensitivity to Miconazole or to any of the excipients. Miconazole Oral gel should not be given concurrently with the drugs tefenadine, astemizole, mizolastine, cisapride.

Side effects : Occasionally, nausea and vomiting have been reported, and with long term treatment, diarrhea. In rare instances, allergic reactions have been reported.

Precaution : Particularly in infants and young children, Repeated use of small Doses is safe caution is required to ensure that the gel does not obstruct the throat. Hence, the gel should not be applied to the back of the tongue and the full dose should be divided into smaller portions.

Micoderm Oral Gel

Pregnancy and Lactation : Micoderm Oral gel should be avoided in pregnant women if possible. The potential hazards should be balanced against the possible benefits. It is not known whether Miconazole is excreted in human milk. Caution should be exercised when prescribing Miconazole Oral Gel to nursing mothers. Pregnancy Category-**N**, Not Classified.

Packing : Micoderm Oral Gel : 15gm in a tube.