

Composition : Terbinafine 1%.

Indications : It is used to treat fungal infections of the skin caused by Trichophyton (e.g. T. Rubrum, T. Mentagrophytes, T. Verrucosum, T. Violaceum), Microsporum canis and Epidermophyton floccosum. It is also used to treat yeast infections of the skin, principally those caused by the genus Candida (e.g. C. albicans).

Dosage and administration : Adult: It can be applied once or twice daily after drying the affected areas thoroughly. Apply the cream to the affected skin and surrounding area in a thin layer and rub in lightly. In the case of intertriginous infections (submammary, interdigital, intergluteal, inguinal) the application may be covered with a gauze strip, especially at night. The likely durations of treatment are as follows. Tinea corporis, cruris: 1 to 2 weeks, Tinea pedis: 1 week, Cutaneous candidiasis: 2 weeks, Pityriasis versicolor: 2 weeks. **Children:** The experience with topical cream in children is still limited and its use cannot therefore be recommended. Or as directed by the registered physician.

Contraindication : It is contraindicated in those patients with hypersensitivity to terbinafine or any of the excipients contained in the cream.

Use in pregnancy and lactation :
Pregnancy: There is no clinical experience with this cream in pregnant women. Therefore, unless the potential

Mycofree Cream

benefits outweigh any potential risk, this cream should not be administered during pregnancy. **Lactation:** Terbinafine is excreted in breast milk and therefore mothers should not receive this cream whilst breast-feeding.

Side effects : Redness, itching or stinging occasionally occur at the site of application. However, treatment rarely has to be discontinued for this reason.

Precautions : It is for external use only. Patient should be avoided contact with eyes.

Packing : Mycofree Cream : 15gm in a tube.