

Sliptin Tablet

Composition : Sitagliptin 25mg, 50mg & 100mg Tablet.

Indications : Sitagliptin is indicated for the treatment of patients suffering from type-II diabetes mellitus. It is also used in combination with metformin or other glycemic control drugs.

Dosage and administration : The recommended dose of Sitagliptin is 100mg once daily. Sitagliptin can be taken with or without food. For patients with mild renal insufficiency (creatinine clearance $\text{crcl} \geq 50\text{ml/min}$), no dosage adjustment is required. For patients with moderate renal insufficiency ($\text{crcl} \geq 30$ to $<50\text{ ml/min}$), the dose of Sitagliptin is 50mg once daily. For patients with severe renal insufficiency ($\text{CrCl} < 30\text{ ml/min}$) or with end-stage renal disease (ESRD) requiring hemodialysis or peritoneal dialysis, the dose of Sitagliptin is 25mg once daily. Or, as directed by the registered physicians.

Contraindication : It is contraindicated in patients who have hypersensitivity.

Side effects : Adverse reactions like upper respiratory tract infection, nasopharyngitis, and headache can occur. Hypoglycemia occurs in patients treated with the combination of Sitagliptin and sulfonylurea.

Use in pregnancy and lactation : Sitagliptin has been assigned to

pregnancy category B by the FDA.

Drug interactions : There was a slight increase in the area under the curve and mean peak drug concentration of Digoxin with the co-administration of Sitagliptin.

Precautions : Dosage adjustment is recommended in patients with moderate or severe renal insufficiency. When used with a sulfonylurea, a lower dose of sulfonylurea may be required to reduce the risk of hypoglycemia.

Packing :

Sliptin 25 : 2x 14's tablets in blister pack.

Sliptin 50 : 2x 14's tablets in blister pack.

Sliptin 100 : 1x 14's tablets in blister pack.