

Composition : Tiemonium Methylsulfate 50mg Tablet & 5mg/2ml Injection.

Indications : It is indicated for treatment of gastroenteritis, diarrhoea, dyscentery, biliary, colic, enterocolitis, cholecystitis, colonopathies, mild cystitis. Visonium is the drug of choice for the treatment of spasmodic dysmenorrhoea.

Dosage and administration : Visonium Tablet : In adults, the usual dosage is 2 to 6 tablets is given in a dose of 100 to 300mg daily in divided doses by mouth. Visonium Injection (IM/IV) : The usual dosage is 5mg (1 ampoule) three times daily by intramuscular or slow intravenous injection. Or, as directed by the registered physician.

Contraindications : This drug should not be used in the following cases :

Glaucoma, Disorders of prostate or bladder, Hyperthyroidism, Paralytic ileus, Pyloric stenosis, Tachycardia, Pulmonary oedema.

Precautions : The drug should be used cautiously in the following situations : Chronic bronchitis, coronary insufficiency, severe renal & hepatic insufficiency. To avoid the risk of hypotension and tachycardia, the injection should be administered slowly.

Drug interaction : Not Known.

Visonium

Tablet & IV/IM Injection

Side effect : Not Known.

Use in pregnancy and lactation : The drug should be used with caution during pregnancy and lactation.

Packing : Visonium Tablet : 5x10's tablets in blister pack.

Visonium Injection (IM/IV) : 1x5's ampoules (2ml) in a blister pack.