

Composition : Aceclofenac 100mg Tablet.

Indications : **Nofenac** is used for the relief of pain and inflammation in the following conditions:

- * Acute and chronic rheumatoid arthritis
- * Osteoarthritis
- * Ankylosing spondylitis
- * Pain and inflammation in orthopaedic, gynecologic, dental and other minor surgery
- * Post operative pain
- * Low back pain and other acute musculoskeletal disorder
- * Primary dysmenorrhea
- * Severe traumatic injuries
- * Severe migraine attacks
- * Renal and biliary colic

Dosage & administration : Adults : The recommended dose is 100mg, twice daily. Children : There are no clinical data on the use of Aceclofenac in children. Elderly : Generally no dose adjustment is necessary. Or as directed by registered physician.

Contraindications : It is contraindicated in patients with known hypersensitivity to Aceclofenac or any of the excipients. It is also contraindicated in patients in whom peptic ulcer or gastrointestinal bleeding, asthma, urticaria, or other sensitivity reaction is precipitated by aspirin or other NSAID's.

Precautions : Symptoms or history of gastrointestinal disease, impaired hepatic, cardiac or renal function or in

Nofenac

Tablet

elderly patients.

Side effects : Side effects to Aceclofenac usually mild and transient. side effects include Gastrointestinal discomfort, nausea, diarrhoea, tinnitus, depression, drowsiness .

Use in pregnancy and lactation :

There are no well controlled studies the safe use of drug in pregnant woman. This drug should not be used during pregnancy unless the benefits to the mother justify the potential risk to the fetus.

Drug interactions : Aceclofenac may exhibit interaction with digoxin, lithium, methotrexate, salicylate, cyclosporine, warfarin, ciprofloxacin, antacids, diuretics, tolbutamide.

Packing : Nofenac : 5 x 10's tablets .