

Composition : Meclizine hydrochloride 50mg Tablet.

Indications : This tablet is indicated for the management of nausea and vomiting, and dizziness associated with motion sickness, vertigo associated with diseases affecting the vestibular system.

Dosage and administration : For the prevention and treatment of nausea, vomiting and vertigo associated with motion sickness, the usual oral dosage of meclizine hydrochloride in adults and children 12 years of age or older is 25-50mg once daily. Vertigo : for the control of vertigo associated with diseases affecting the vestibular system, the recommended dose is 25 to 100mg daily, in divided doses, depending upon clinical response. Motion Sickness : The initial dose of 25 to 50mg should be taken one hour prior to embarkation for protection against motion sickness. Thereafter, the dose may be repeated every 24 hours for the duration of the journey.

Contraindication : Meclizine hydrochloride is contraindicated in individuals who have shown a previous hypersensitivity to it.

Side effects : Drowsiness, dry mouth and, on rare occasions, blurred vision have been reported.

Precautions : Due to its potential anticholinergic action, this drug should be used with caution in patients with asthma, glaucoma, or enlargement of the prostate gland.

Nomasic Tablet

Use in pregnancy and lactation : Pregnancy Category **B** by FDA.

Lactation : Safety for use in the nursing mother has not been established.

Drug interactions : Additive CNS depression may occur where meclizine hydrochloride are administered concomitantly with other CNS depressants including barbiturates, tranquilizers and alcohol.

Packing : Nomasic Tablet : 5x10's tablets in blister pack.