

**Composition :** Diclofenac Potassium 25mg & 50mg Tablet.

**Indications :** It is indicated for the treatment of pain and inflammation in the following diseases- # Rheumatoid arthritis, # Osteoarthritis, # Ankylosing spondylitis, # Acute gout, # Migraine attacks, #Acute musculoskeletal disorders and trauma such as peri-arthritis (especially frozen shoulder), tendinitis, tenosynovitis, bursitis, sprains, strains, fracture, low back pain, # Orthopaedic, dental and other minor surgery.

**Dosage and administration :** Adults : Generally for adults 100- 150mg. daily in 2-3 divided doses.

In milder cases, as well as for children over 14 years 75-100mg. daily in 2-3 divided doses.

In migraine an initial dose of 50mg should be taken at the first signs of an impending attack. In cases where relief 2 hours after the first dose is not sufficient, a further dose of 50mg may be taken. If needed, further doses of 50mg may be taken at intervals 4-6 hours, not exceeding a total dose of 200mg per day .or as directed by the registered physician.

**Side effects :**Gastro intestinal discomfort, nausea, diarrhoea and rare G.I. bleeding and ulceration.

**Precautions :** Symptoms or history of gastro intestinal disease, impaired

## Nopain Tablet

hepatic, cardiac or renal function.

**Use in pregnancy and lactation :** Pregnancy Category C. There are no well controlled studies in pregnant woman. This drug should not be used during pregnancy unless the benefits to the mother justify the potential risk to the fetus.

**Packing :**

**Nopain-25 :** 10 x 10's tablets.

**Nopain-50 :** 5 x 10's tablets.