

**Composition :** Ofloxacin 200mg & 400mg Tablet.

**Indications :** The following infections caused by Ofloxacin susceptible Staphylococcus, Streptococcus pyogenes, Hemolytic streptococci, Streptococcus pneumoniae, Peptostreptococcus sp., Neisseria gonorrhoeae, Escherichia coli, Citrobacter sp., Shigella sp., Klebsiella pneumoniae, Enterobacter sp., Serratia sp., Proteus sp., Pseudomonas aeruginosa, Haemophilus influenzae, Acinetobacter sp., and Campylobacter sp.,

- a) Pneumonia, Chronic Bronchitis, Diffuse panbronchiolitis, Bronchiectasis with infection, Secondary infections of chronic respiratory diseases.
- b) Pharyngolaryngitis, Tonsillitis and Acute bronchitis.
- c) Pyelonephritis, Cystitis, Prostatitis, Epididymitis, Gonococcal urethritis, Non-gonococcal urethritis.
- d) Intrauterine infection, Adnexitis, Bartholinitis.
- e) Folliculitis, Furuncle, Furunculosis, Carbuncle, Erysipelas phlegmon, Lymphangitis, Felon, Subcutaneous abscess, Spiradenitis, Acne conglobata, Infectious atheroma, Perianal abscess.
- f) Mastadenitis, Superficial secondary infections after traumas, Burns, Surgery, Traumas.
- g) Cholecystitis, Cholangitis.
- h) Otitis media, Sinusitis.
- i) Blepharitis, Hordeolum, Dacryocystitis, Tarsadenitis, Keratohelcosis.
- j) Bacillary dysentery, Enteritis.
- k) Periodontitis, Pericoronitis, Gnathitis.
- l) Skin & soft tissue infections.

**Dosage and administration :** General dosage recommendation : The dose of OFLACIN is determined by the type and severity of the infection. The dosage range of adults is 200mg

## Oflacin Tablet

to 800mg daily. Up to 400mg may be given as a single dose, preferably in the morning, larger doses should be given as two divided doses. OFLACIN should not be taken within two hours of magnesium/aluminium containing antacids or iron preparations since reduction of absorption of ofloxacin can occur.

**Lower Urinary Tract Infections :** 200-400mg daily preferably in the morning.

**Upper Urinary Tract Infections :** 200-400mg daily preferably in the morning, increasing, if necessary, to 400mg twice daily.

**Lower Respiratory Tract Infections :** 400mg daily preferably in the morning increasing, if necessary, to 400mg twice daily.

**Uncomplicated Urethral and Cervical Gonorrhoea :** A single dose of 400mg.

**Non Gonococcal Urethritis and Cervicitis :** 400mg daily in single or divided doses.

**Impaired renal function :** 100mg should be given every 48 hours, when creatinine clearance is less than 20ml/minute (serum creatinine greater than 5mg/dl).

**Elderly :** No adjustment of dosage is required in the elderly.

**Duration of treatment :** The usual treatment period is 5-10 days except in uncomplicated gonorrhoea where a single dose is recommended or as directed by a registered physician.

**Use in pregnancy and lactation :** Ofloxacin has been assigned to pregnancy category **C** by the FDA.

**Contraindication :** Patients with hypersensitivity to Ofloxacin.

**Packing :**

**Oflacin 200 :** 2x10's tablets in blister pack.

**Oflacin 400 :** 2x10's tablets in blister pack.