

Composition : Domperidone 10mg Tablet & 5mg/5ml Suspension.

Indications : Paridon is indicated for the treatment of

1. Functional dyspepsia 2. Stimulation of gut mobility (a) non-ulcer dyspepsia (b) esophageal reflux, reflux esophagitis and gastritis 3. Acute treatment of nausea and vomiting, in adults. 4. For up to 12 weeks treatment of nausea and vomiting caused by L-dopa and bromocriptine.

Dosage & administration : The recommended oral dose of Paridon Suspension for :

Adults : 10 - 20ml every 4-8 hours daily.

Children : 0.2 - 0.4mg/kg every 4 - 8 hours daily.

For the treatment of nausea and vomiting - Adults (including the elderly) : 10-20 mg orally at 4-8 hourly intervals. Or, as directed by the physician.

For the treatment of symptoms of functional dyspepsia & others : Adults (including the elderly): Tablets up to 10-20mg orally 3 times daily before meals and 10-20mg at night depending on clinical response. A course of treatment should not exceed 12 weeks. Or, as directed by the physician.

Contraindication : Domperidone is contraindicated to patients who have known hypersensitivity to this drug.

Precautions : Domperidone should be used with absolute caution in case of children because there may be an increased risk of extra pyramidal reactions in young children because of

Paridon

Tablet & Suspension

an incompletely developed blood brain barrier.

Side effects : Domperidone produces a rise in serum prolactin which may be associated with galactorrhoea and less frequently with gynaecomastia, breast enlargement or soreness, there have been reports of reduced libido. Occasional rashes and other allergic phenomena have been reported.

Drug interactions : Domperidone may reduce the hypoprolactinemic effect of bromocriptine. The actions of Domperidone on gastrointestinal function may be antagonized by antimuscarinics and opioid analgesics. Domperidone may enhance the absorption of concomitantly administered drugs, particularly in patients with delayed gastric emptying.

Use in pregnancy and lactation : Pregnancy Category **C** by FDA.

Packing : Paridon : 7x14's tablets in blister pack.

Paridon suspension : 100ml bottle.

