

**Composition :** Betamethasone (as 17 valerate) 0.05%, □ Phenylephrine hydrochloride 0.1% & □ Lignocaine hydrochloride 2.5%.

**Indications :** Relitch is indicated for: relief of the symptoms of itching, irritation, discomfort or pain associated with local non-infective anal or peri-anal conditions such as external hemorrhoids.

**Dosage and administration :** a course of treatment should be limited to 7 days. Apply a small amount of ointment two or three times a day initially, using the applicator if internal administration is required. When inflammation is subsiding, once-daily application is sufficient in most cases. Or as directed by the physician

**Contraindication :** Primary cutaneous viral infections (e. g. herpes simplex, chickenpox). Hypersensitivity to any components of the drug. The use of this ointment is not indicated in the treatment of primarily infected skin lesions caused by infection with fungi (e. g. candidiasis, tinea); or bacteria (e. g. impetigo).

**Pregnancy and lactation :** There is inadequate evidence of safety in human pregnancy.

**Side effects :** Signs of hypersensitivity may appear. Prolonged and intensive treatment with highly active corticosteroid preparations may cause local atrophic changes in the skin.

## Relitch Ointment

**Precautions :** Long-term continuous therapy should be avoided where possible, particularly in infants and children, as adrenal suppression can occur.

**Packing :** Relitch rectal ointment : 15 gram in a tube.