

## Rofumil Tablet

**Composition :** Roflumilast 500mcg Tablet.

**Indications :** Roflumilast is indicated for the maintenance treatment of severe COPD associated with chronic bronchitis with a history of frequent exacerbation.

**Dosage and administration :** The recommended dose of Roflumilast is one 500 microgram(mcg) tablet once daily. The tablet should be swallowed with water and can be taken with or without food. Or, as directed by the registered physician.

**Contraindication :** It is contraindicated in patients with known hypersensitivity to Roflumilast or any other component of the medicine.

**Side effects :** The most common reported side effects are weight decrease, anorexia, insomnia, nausea, headache, GI upset.

**Precautions :** Roflumilast is not a bronchodilator and therefore is not indicated for the relief of acute symptoms. It should not be given in latent infections an psychiatric symptoms or taking other medications associated with psychiatric reactions. Patients are advised to report any changes in behavior or mood.

**Use in pregnancy and lactation :** Roflumilast is pregnancy category C by the FDA.

**Drug interactions :** It shows drug interactions with cimetidine and other inhibitors of CYP3A4 (e.g. erythromycin, ketoconazole) and strong P450 inducers (e.g. rifampicin, carbamazepine,

phenobarbital, phenytoin).

**Packing : Rofumil :** 2x10's tablets in blister pack.