

Composition : Zinc Sulfate Monohydrate Eq. to 10mg of elemental Zinc/5ml Syrup.

Indications : It is indicated for the following conditions-growth retardation, RTI, alopecia, dermatitis, diarrhoea, immunological dysfunction, psychological disturbances, gonadal atrophy, impaired spermatogenesis, congenital malformation.

Dosage and administration : Children under 10kg : 22.5mg ($2\frac{1}{4}$ spoonful) Ziton syrup daily in divided doses. Children over 10kg to 30kg : 22.5mg ($2\frac{1}{4}$ spoonful) 1 to 3 times daily. Adult & children over 30kg : 45mg ($4\frac{1}{2}$ spoonful) 1 to 3 times daily after food.

Or, as directed by the registered physician.

Side effects : Prolonged use may lead to copper deficiency and anemia which has responded to withdrawal of zinc and symptomatic therapy, zinc sulfate is the form of zinc often used for oral administration causes adverse gastrointestinal effects.

Precautions : Concurrent administration of zinc salt with penicillamine might diminish the effect of penicillamine. The absorption of zinc, although poor, may be decreased by various compounds including some foods. Chelation may occur with tetracyclines.

Ziton Syrup

Packing : Ziton :100ml in amber glass bottle.