

**Composition :** Each Metered dose (Puff) contains Salbutamol 100 mcg as Salbutamol Sulphate.

**Indication : Bronchospasm :** Ventil is indicated for the treatment of prevention of bronchospasm in patients 4 years of age and older with reversible obstructive airway disease.

**Exercise-induced Bronchospasm :** Ventil is indicated for the prevention of exercise-induced bronchospasm in patients 4 years of age and older.

**Dosage and administration :** For treatment of acute episodes of bronchospasm or prevention of asthmatic symptoms, the usual dosage for adults and children 4 years of age and older is 2 inhalations repeated every 4 to 6 hours; in some patients, 1 inhalation every 4 hours may be sufficient. More frequent administration or a larger number of inhalations is not recommended. **Exercise-Induced Bronchospasm Prevention:** The usual dosage for adults and children 4 years and older is 2 inhalations 15 to 30 minutes before exercise. Or, as directed by the registered physician.

**Side effects :** Mild tremor and headache have been rarely reported. These usually disappear with continuous treatment. There have been very rare reports of transient muscle cramp. Hypersensitivity reactions including angioedema, urticaria, bronchospasm, hypotension and collapse have been reported very rarely. As with other inhalation therapy, the potential for paradoxical bronchospasm should be kept in mind. If

## Ventil

HFA Inhaler  
200 Puffs

it occurs, the preparation should be discontinued immediately and alternative therapy should be instituted.

**Contraindication :** Salbutamol is contraindicated in patients with a history of hypersensitivity to Salbutamol or any other components of Salbutamol.

**Use in pregnancy and lactation : Pregnancy category C.** There are no adequate and well- Controlled studies of Salbutamol sulphate in pregnant women. Salbutamol inhaler should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. It is not known whether the components of Salbutamol excreted in human milk.

**Drug interactions :** Salbutamol and nonselective b-blocking drugs eg, propranolol, should not be prescribed together.

**Precautions :** Salbutamol as with all sympathomimetic amines should be used with caution in patients with cardiovascular disorders, especially coronary insufficiency, hypertension and cardiac arrhythmia; in patients with convulsive disorders, hyperthyroidism, or diabetes mellitus.

**Packing : Ventil HFA Inhaler :** Each box contains one inhaler of 200 Puffs.