

Composition : Tobramycin + Dexamethasone (3mg+1mg)/ml.

Indications : Texa is indicated for steroid-responsive inflammatory ocular conditions for which a corticosteroid is indicated and where superficial bacterial ocular infection or a risk of bacterial infection exists.

Dosage and administration : One or two drops instilled into the conjunctival sac(s) every four to six hours. During the initial 24 to 48 hours, the dosage may be increased to one or two drops every two (2) hours. Frequency should be decreased gradually as warranted by improvement in clinical signs. Or, as directed by the registered physician.

Contraindication : Epithelial herpes simplex keratitis (dendritic keratitis), vaccinia, varicella, and many other viral diseases of the cornea and conjunctiva. Mycobacterial infection of the eye. Fungal diseases of ocular structures. Hypersensitivity to a component of the medication.

Precautions : As with other antibiotic preparations, prolonged use may result in overgrowth of nonsusceptible organisms, including fungi.

Side effects : The most frequent adverse reactions to topical ocular Tobramycin are hypersensitivity and localized ocular toxicity, including lid itching and swelling, and conjunctival erythema.

Texa

Sterile Ophthalmic
Suspension

Drug Interaction : No information provided.

Pregnancy and lactation : There are no adequate and well controlled studies in pregnant women. Texa suspension should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Packing : 5ml in plastic dropper bottle.