

Trialon Nasal Spray

Composition : Trialon Nasal Spray : Each metered spray contains Triamcinolone Acetonide BP 55 mcg.

Indications : This nasal spray indicated for the treatment of the nasal symptoms of seasonal and perennial allergic rhinitis in adults and children 6 years of age and older.

Dosage and administration : Adults and children 12 years of age and older: The recommended starting and maximum dose is 220 mcg per day as two sprays in each nostril once daily. Children 6 to 12 years of age: The recommended starting dose is 110 mcg per day given as one spray in each nostril once daily. The maximum recommended dose is 220 mcg per day as two sprays per nostril once daily. Children 2 to 5 years of age : Starting and maximum dose 110 mcg per day as one spray in each nostril once daily. Not recommended for children under 2 years of age. Or, as directed by the registered physician.

Side effects : Generally, Triamcinolone Acetonide, is well tolerated. The few side effects such as: pharyngitis, epistaxis, increase in cough, headache, infection, sinusitis and vomiting may occur.

Contraindication : It is contraindicated in patients with a hypersensitivity to Triamcinolone Acetonide or any other ingredients in this preparation.

Use in pregnancy and lactation : Pregnancy category C. There are no adequate and well controlled studies in pregnant women. Triamcinolone Acetonide nasal spray, like other corticosteroids, should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

It is not known whether Triamcinolone Acetonide is excreted in human milk. Because other corticosteroids are excreted in human milk, caution should be exercised when Triamcinolone Acetonide nasal spray is administered to a nursing woman.

Drug interactions : No significant drug interactions has been reported.

Precautions : It should be used with caution in patients with active or quiescent tuberculous infection of the respiratory tract or in patients with untreated fungal, bacterial, or systemic viral infections or ocular herpes simplex. In clinical studies with Triamcinolone Acetonide nasal spray, the development of localized infections of the nose and pharynx with *Candida albicans* has rarely occurred.

Overdose : Like any other nasally administered corticosteroids, acute overdosing is unlikely in view of the total amount of active ingredient present. In the event that the entire contents of the bottle were administered all at once, via oral or nasal application, clinically significant adverse events would most likely not result. The patient may experience some gastrointestinal upset.

Storage : Store below 25°C, protect from light.

Packing : Trialon Nasal Spray : Each bottle contains 120 metered doses for sprays.