

Composition : Triamcinolone Acetonide 0.1%.

Indications : It is indicated for aphthous ulcers, ulcerative stomatitis, denture stomatitis, desquamative gingivitis, erosive lichen planus and lesions of the oral mucosa of traumatic origin.

Dose & Administration : Adults and Children : To be applied to the lesion at bedtime and two to four times daily, preferably after meals. Apply to the affected area; do not rub in.

Elderly : Corticosteroids should be used sparingly and for short periods of time. or as directed by the registered physician.

Contraindication : This medicine is contraindicated to patients with hypersensitivity to any component of this product.

Precautions : Adrenal suppression can occur with prolonged use of topical corticosteroids or with occlusion. These effects are more likely to occur in infants and children and courses of treatment in childhood should be limited to 5 days. Dentures may act in the same way as an occlusive dressing.

Side effects : Triamcinolone acetonide is well tolerated. Where adverse reactions occur they are usually reversible on cessation of therapy. The possibility of the systemic effects which are associated with all steroid therapy should be considered.

Trialon **Oral Paste**

Use in pregnancy and lactation : Triamcinolone should only be prescribed when the benefits to the mother & child outweigh the risks. Some of corticosteroids may pass into breast milk, but no data are available for Triamcinolone Acetonide. Pregnancy Category **C**.

Drug interaction : Barbiturates, phenytoin, rifampicin, rifabutin, carbamazepine, primidone and aminoglutethimide may enhance the metabolic clearance of corticosteroids resulting in decreased therapeutic effects.

Packing : Trialon Oral Paste : 10 gram in a tube.