

Composition : D-CAP 800 : Each Soft Gelatin Capsule Contains Colecalciferol (Vitamin D₃) BP 800 IU.

D-CAP 1000 : Each Soft Gelatin Capsule Contains Colecalciferol (Vitamin D₃) BP 1000 IU.

D-CAP 2000 : Each Soft Gelatin Capsule Contains Colecalciferol (Vitamin D₃) BP 2000 IU.

D-Cap 20000 : Each Soft Gelatin Capsule Contains Colecalciferol (Vitamin D₃) USP 20000 IU.

D-Cap 40000 : Each Soft Gelatin Capsule Contains Colecalciferol (Vitamin D₃) USP 40000 IU.

Indication : It is indicated in the prevention and treatment of vitamin D₃ deficiency. As an adjunct to specific therapy for osteoporosis in patients with vitamin D₃ deficiency or at risk of vitamin D₃ insufficiency.

Dosage & Administration : Use in adults : The recommended dose for :

Prevention of vitamin D deficiency : 1000 IU/day or (20000 IU) once a month.

Addition to specific therapy for osteoporosis : 1,000 IU/day or (20000 IU) once a month.

Treatment of vitamin D deficiency : 1000 IU-4,000 IU/day for up to 12 weeks, followed by maintenance therapy of 1400 IU-2000 IU/day or 40000 IU once weekly for 7 weeks, followed by maintenance therapy such as 2-3 capsule per month.

Use in children and adolescents : The recommended dose for :

* **Prevention of deficiency 10-18 years :** doses of up to 1000 IU/day may be required to prevent deficiency in some children.

* **Treatment of deficiency 10-18 years :** 2000 IU/day for 6 weeks followed by a maintenance therapy of 400 IU-1000 IU/day (such as one 1000 IU soft capsule per day).

Or, as directed by the registered physician.

Contraindications : It is contraindicated in patients with known hypersensitivity to Vitamin D₃ or any other components of this product.

Precautions : Vitamin D₃ 20000 IU & Vitamin D₃ 40000 IU Capsules should be used with caution in patients with impaired renal function, impaired renal calcium and phosphate excretion, a tendency to form kidney stones (calculi), hypercalcemia, hypercalciuria.

D-CAP
Capsule



DRUG
INTERNATIONAL
LTD.

Side effects : The side effects of this product are hypercalcaemia, hypercalciuria, pruritus, rash, urticaria, Constipation, flatulence, nausea, abdominal pain, diarrhoea chronic (increased urination, increased thirst, loss of appetite, weight loss, kidney stones, kidney calcification, calcification may in tissues outside the bone) episodes occur.

Use in Pregnancy and Lactation : If anyone are pregnant or breast-feeding, think may be pregnant or are planning to have a baby, should consult with doctor or pharmacist for advice before taking this medicine.

Drug-Interactions : Concomitant treatment with phenytoin or barbiturates, glucocorticoids can decrease the effect of Vitamin D₃. Simultaneous treatment with ion exchange resins such as cholestyramine or laxatives such as paraffin oil may reduce the gastrointestinal absorption of Vitamin D₃. The cytotoxic agent actinomycin and imidazole antifungal agents interfere with Vitamin D₃ activity.

Storage : Store at a cool & dry place, protect from light & moisture. Keep the medicine out of reach of children.

Packing : D-CAP 800 : Each box contains 50's soft capsules in blister pack. □

D-CAP 1000 : Each box contains 50's soft capsules in blister pack. □

D-CAP 2000 : Each box contains 50's soft capsules in blister pack.

D-Cap 20000 : Each box contains 10's soft capsules in blister pack.

D-Cap 40000 : Each box contains 10's soft capsules in blister pack.