

Composition : Nopain-25 : Each Film Coated Tablet Contains: Diclofenac Potassium USP 25mg.

Nopain-50 : Each Film Coated Tablet Contains: Diclofenac Potassium USP 50mg.

Pharmacology : Diclofenac has analgesic, anti-inflammatory and antipyretic properties. The mechanism of action of Diclofenac like that of other NSAIDs, is not completely understood but involves inhibition of cyclooxygenase (COX-1 and COX-2). Diclofenac is 100% absorbed after oral administration. Diclofenac is eliminated through metabolism and subsequent urinary and biliary excretion of the glucuronide and the sulfate conjugates of the metabolites. Little or no free unchanged diclofenac is excreted in the urine.

Indications : It is indicated for the treatment of pain and inflammation in the following diseases-

Rheumatoid arthritis, Osteoarthritis, Ankylosing spondylitis, Acute gout, Migraine attacks, Acute musculoskeletal disorders and trauma such as peri-arthritis (especially frozen shoulder), tendinitis, tenosynovitis, bursitis, sprains, strains, fracture, low back pain, Orthopaedic, dental and other minor surgery.

Dosage and administration : Adults : Generally for adults 100-150mg daily in 2-3 divided doses.

In milder cases, as well as for children over 14 years 75-100mg daily in 2-3 divided doses.

In migraine an initial dose of 50mg should be taken at the first signs of an impending attack. In cases where relief 2 hours after the first dose is not sufficient, a further dose of 50mg may be taken. If needed, further doses of 50mg may be taken at intervals 4-6 hours, not exceeding a total dose of 200mg per day. Or, as directed by the registered physician.

Contra Indications : Peptic ulcer and hypersensitivity to the active substance.

NOPAIN

Tablet



DRUG
INTERNATIONAL
LTD.

Precautions : Symptoms or history of gastro intestinal disease, impaired hepatic, cardiac or renal function or cautions use in elderly patients.

Side effects : Gastro intestinal discomfort, nausea, diarrhoea and rare G.I. bleeding and ulceration.

Use in Pregnancy and lactation : Pregnancy Category C. There are no well controlled studies in pregnant woman. This drug should not be used during pregnancy unless the benefits to the mother justify the potential risk to the fetus.

Use in Child : There is no data available.

Drug Interactions : Diclofenac may interact with the following drugs- other NSAID, diuretics, antihypertensive agents, digoxin, lithium, methotrexate, ciclosporin, quinolone antibiotics and phenytoin.

Overdose : Symptoms of overdoses are headache, nausea, vomiting, epigastric pain.

Storage : Store below 30°C in a dry place.

Packing : Nopain-25 : Each box contains 10 x 10's tablets in blister pack.

Nopain-50 : Each box contains 5 x 10's tablets in blister pack.