

**Composition :** Each film coated tablet contains Tolperisone Hydrochloride INN 50mg.

**Pharmacology :** Tolperisone is a centrally acting muscle relaxant that acts at the reticular formation in the brain stem by blocking voltage-gated sodium and calcium channels.

**Indications :**

1. Treatment of increased tone of skeletal muscles due to organic neurological disorders (injury of the pyramidal tract, multiple sclerosis, myelopathy, encephalomyelitis).
2. Treatment of muscular spasm, muscular contracture, rigidity, spinal automatism.
3. In the treatment of vascular diseases (Obliterative arteriosclerosis, diabetic angiopathy, obliterative thromboangitis, Raynaud's disease, diffuse scleroderma).
4. In individual cases, post-thrombotic venous and lymphatic circulation disorders and crural ulcer.

**Dosage and administration :** The recommended dose is as followings :

Adult : 50-150mg three times daily according to the individual requirement and tolerance of the patient.

Children (from 3 months to 6 years) : 5mg/kg/day in three divided doses.

Children (from 6 to 14 years) : 2-4mg/kg/day in three divided doses.

Or, as directed by the registered physician.

**Contraindication :** It is contraindicated in patients with a history of hypersensitivity to any component of the product, myasthenia gravis and nursing mother.

**Lexaton**

Tablet



DRUG  
INTERNATIONAL  
LTD.

**Precautions :** In case of children, the prescribed dose and duration of treatment should closely be observed.

**Side effects :** Possible side-effects include sometimes muscular weakness, headache, nausea, vomiting and rarely hypersensitivity reactions (itching, redness of the skin, skin rash) may occur.

**Use in pregnancy and lactation :** Lexaton should be used in pregnancy according to physician's advice. It should not be used during breast feeding.

**Use in Child :** There is no data available.

**Drug Interactions :** Concomitant use of methocarbamol with Tolperisone Hydrochloride has been reported to cause disturbance of visual accommodation.

**Overdose :** There is no data available.

**Storage :** Store below 30°C in a dry place.

**Packing :** Each box contains 4 x 14's tablets in blister pack.