

Composition : Each Film Coated Tablet Contains Paracetamol BP 500.00mg and Caffeine BP 65.00mg.

Pharmacology : Paracetamol is a non-opioid, non-salicylate analgesic. The site and mechanism for the analgesic effect of Paracetamol has not been determined but is thought to primarily involve central actions. Paracetamol appears to be widely distributed throughout most body tissues except fat. A relative small portion (20%) of Paracetamol is bound to plasma protein. Paracetamol is eliminated from the body primarily by formation of glucuronide and sulfate conjugates in a dose dependent manner. Paracetamol is primarily metabolized in the liver by first-order kinetics and involves three principal separate pathways. Less than 9% of Paracetamol is excreted unchanged in the urine.

Indications : It is indicated for the treatment of most painful and febrile conditions, for example, headache, including migraine, backache, toothache, rheumatic pain and dysmenorrhea, and relief of the symptoms of colds, influenza and sore throat.

Dosage and administration : Adults and children 16 years and over. Two tablets every 4-6 hours up to four times daily. Do not exceed 8 tablets in 24 hours. Children and children aged 12-15 years: One tablet every 4-6 hours up to four times daily. Do not exceed 4 tablets in 24 hours. Not recommended for children under 12 years. Or, as directed by the registered physician.

Contraindication : It is contraindicated in patients with known hypersensitivity to active ingredient or any other components of this product.

Precautions : Caution should be advised in the administration of paracetamol to patients with renal or hepatic impairment. Excessive intake of caffeine (e.g. coffee, tea and some canned drinks) should be avoided while taking this product. Patients should be advised to consult their doctor if their

Momodol Extra Tablet



headaches become persistent. Patients should be advised not to take other paracetamol-containing products concurrently.

Side effects : Side effects include : Thrombocytopenia, Agranulocytosis, Anaphylaxis, Cutaneous hypersensitivity reactions including skin rashes, angioedema and Stevens Johnson syndrome, toxic epidermal necrolysis, Bronchospasm, Hepatic dysfunction, Nervousness, Dizziness.

Use in pregnancy and lactation : Paracetamol-caffeine is not recommended for use during pregnancy due to the possible increase risk of lower birth weight and spontaneous abortion associated with caffeine consumption, Caffeine in breast milk may potentially have a stimulating effect on breast fed infants. Due to the caffeine content of this product it should not be used in pregnant or breast feeding women.

Use in Child : There is no data available.

Drug interactions : The speed of absorption of paracetamol may be increased by metoclopramide or domperidone and absorption reduced by colestyramine. The anticoagulant effect of warfarin and other coumarins may be enhanced by prolonged regular daily use of paracetamol with increased risk of bleeding, occasional doses have no significant effect.

Overdose : There is no data available.

Storage : Store below 30°C in a dry place.

Packing : Each box contains 60's tablets in blister pack.