

Composition : Each film coated tablet contains Micronised purified flavonoid fraction 500mg (Diosmin BP 450mg and Hesperidin BP 50mg).

Pharmacology : Diosmin and Hesperidine is a phlebotonic drug and a vascular protecting agent. It reinforces venous tone by prolonging the activity of parietal noradrenaline. Thus Diosmin and Hesperidine decreases venous capacitance, venous distensibility and venous emptying time. Diosmin and Hesperidine protects the microcirculation by fighting the microcirculation damaging process. It combats venous inflammation by decreasing leukocyte activation and as consequence by inhibiting the release of inflammatory mediators, principally free radicals and prostaglandin. Thus Diosmin and Hesperidine normalizes capillary permeability and strengthens capillary resistance.

Indications : This medicine is venotonic (it increases venous tone) and a vasculoprotector (it increases resistance in small blood vessels). It is recommended for treating venous circulation disorders (swelling leg, pain, restless legs) and for treating symptoms due to acute hemorrhoidal attack.

Dosage and administration : **Venous Insufficiency:** 2 tablets daily, one at midday and one at night upto 2 months. **Acute hemorrhoidal attack:** 6 tablets daily for the first 4 days followed by 4 tablets daily for the next 3 days. **Chronic hemorrhoidal attack:** 2 tablets daily. The tablets should be taken at meal times. Or, as directed by the registered physician.

Contraindication : It is contraindicated in patients with known hypersensitivity to diosmin, hesperidin or any other components of this drug.

Ardium

Tablet



**DRUG
INTERNATIONAL
LTD.**

Precaution : This medicine is not a substitute for a specific treatment of other anal disorders. Avoid long-term use. If the symptoms persist, consult your doctor.

Side effects : The most common side effects are mild stomach discomforts and neurovegetative disorders such as hypersomnia, fatigue and increased appetite.

Use in pregnancy and lactation : There are no adequate and well-controlled studies in pregnant women. This drug should be used during pregnancy only if clearly needed. Breastfeeding is not recommended during the treatment period.

Use in Child : The safety and effectiveness in children have not been established.

Drug Interactions : Metabolism of metronidazole may be reduced by concurrent admin of diosmin-hesperidin.

Overdose : No overdosage symptoms have been observed.

Storage : Store at a cool and dry place, protected from light and moisture. Keep out of reach of children.

Packing : Each box contains 3 x 10's tablets in blister pack.