

**Composition :** Each tablet contains 5mg Vinpocetin USP.

**Pharmacology :** **Camiton** improves cerebral metabolism : Increases the glucose and oxygen consumption and utilization of the brain tissue; improves the hypoxia tolerance of the brain cells; shifts glucose metabolism toward the energetically more favorable aerobic pathway, exerts antioxidant activity; as a result of all these effects it is a cerebroprotective compound.

**Camiton** considerably improves cerebral microcirculation :

Inhibits platelet aggregation : reduces pathologically increased blood viscosity; promotes the oxygen transport into the brain tissue.

**Camiton** selectively and intensively increases cerebral blood flow and the cerebral fraction of cardiac output, reduces cerebral vascular resistance without affecting systemic circulation.

**Indications :** All forms of acute and chronic cerebral circulatory insufficiency: TIA, reversible ischaemic neurological deficiency, progressive stroke, completed stroke, multi-infarct dementia, cerebral arteriosclerosis, hypertensive encephalopathy etc. For the reduction of psychic or neurological symptoms of cerebral insufficiency (e.g. memory disturbances, dizziness, headache, aphasia, locomotor disorders etc.).

**Ophthalmology :** It can be used for the treatment of vascular disorders of the chorioid and retina due to arteriosclerosis, vasospasm, macula degeneration, arterial or venous thrombosis or embolism, and glaucoma secondary to the above mentioned disorders.

**CAMITON**

Tablet



DRUG  
INTERNATIONAL  
LTD.

**Otology :** For the treatment of impaired hearing of vascular or toxic (iatrogenic) origin, presbycusis, Menier's disease, cochleovestibular neuritis, tinnitus and dizziness of labyrinth origin.

**Dosage & Administration :** 1-2 tablet 3 times daily. Or, as directed by the registered physician.

**Side effects :** Transient hypotension, tachycardia may occur.

**Contraindication :** Pregnancy or hypersensitivity to any component of the tablet.

**Precautions :** Vinpocetine might increase the risk of miscarriage and might cause harm to the fetus. Avoid using.

**Drug interaction :** Incompatible with Heparin.

**Use in pregnancy and lactation :** Vinpocetine is POSSIBLY UNSAFE when used in pregnant women or those who may become pregnant.

**Use in Child :** Safety and effectiveness in children have not been established.

**Overdose :** There is no data available.

**Storage :** Store below 30°C in a dry place.

**Packing :** Each box contains 7 x 14's tablets in blister pack.