

**Composition :** Each ml ophthalmic solution contains Bromfenac 0.9mg as Bromfenac sodium INN.

**Indications :** Bromfenac ophthalmic solution is indicated for treatment of postoperative inflammation and in patients who have undergone cataract extraction, prevention of miosis in cataract surgery, treatment of pseudophakic cystoid macular edema (CME), treatment of external anterior ocular inflammatory diseases such as blepharitis, conjunctivitis and scleritis.

**Dosage and administration :** For the treatment of postoperative inflammation and in patients who have undergone cataract extraction, one drop of Bromfenac ophthalmic solution should be applied to the affected eye (s) two times daily beginning 24 hours after cataract surgery and continuing through the first two weeks of the postoperative period. Prevention of miosis in cataract surgery: beginning 1 hour prior to surgery, instill bromfenac every 30 minutes. Treatment of pseudophakic CME: 1 drop 2 times daily for 3 months. Or, as directed by the registered physician.

**Contraindication :** Bromfenac ophthalmic solution is contraindicated in patients with known hypersensitivity to any ingredient in the formulation.

**Side effects :** The most commonly reported adverse experiences following use of bromfenac include abnormal sensation in eye, conjunctival hyperemia, eye irritation (including burning/stinging), eye pain, eye pruritus, eye redness, headache and iritis.

**Precautions :** Use of topical nonsteroidal anti-inflammatory drugs (NSAIDs) in patients

**Eyebrom**

Sterile Eye Drops



**DRUG  
INTERNATIONAL  
LTD.**

with complicated ocular surgeries, corneal denervation, corneal epithelial defects, diabetes mellitus, ocular surface diseases (e.g. dry eye syndrome), rheumatoid arthritis or repeat ocular surgeries within a short period of time may be at increased risk for corneal adverse events which may become sight threatening. Topical NSAIDs should be used with caution in patients with known bleeding tendencies or who are receiving other medications, which may prolong bleeding time. Bromfenac ophthalmic solution should not be administered while wearing contact lenses.

**Use in Pregnancy and lactation :** There are no adequate and well controlled studies in pregnant women. This drug should be used during pregnancy only if the potential benefit to the mother justifies the potential risk to the fetus. Caution should be exercised when bromfenac ophthalmic solution is administered to a nursing women.

**Drug interactions :** Not known.

**Storage :** Store at a cool and dry place, protected from light and moisture. Keep out of reach of children. Do not use longer than one month after first opening.

**Packing :** **Eyebrom** Eye Drops : 5ml plastic dropper bottle.