

**Composition : Dialiptin-50** : Each film coated tablet contains Vildagliptin INN 50mg.

**Pharmacology** : Vildagliptin is a dipeptidylpeptidase-4 (DPP-4) inhibitor, which is believed to exert its actions in patients with type 2 diabetes by slowing the inactivation of incretin hormones. Vildagliptin increases insulin release and decreases glucagon levels in the circulation in a glucose-dependent manner.

**Indications** : Vildagliptin is indicated for the treatment of type 2 diabetes mellitus (in combination with metformin or with a sulphonylurea or with a thiazolidinedione).

**Dosage and administration : Adult over 18 years:** Vildagliptin administered in combination with metformin or a thiazolidinedione, 50mg twice daily; in combination with a sulphonylurea, 50mg daily in the morning. Or, as directed by the registered physician.

**Contraindication** : It is contraindicated for patients with diabetic ketoacidosis, hepatic impairment, kidney dysfunction.

**Precautions** : Caution should be employed for elderly patients. It should be used with caution in patients with impaired hepatic function and heart failure.

**Side effects** : The most common side effects associated with Vildagliptin therapy include headache, tremor, dizziness, nausea and asthenia. Less commonly constipation, hypoglycemia.

**Dialiptin-50**

Tablet



DRUG  
INTERNATIONAL  
LTD.

**Pregnancy and lactation** : Vildagliptin should be avoided during pregnancy or breast feeding unless essential.

**Use in Child** : Safety and effectiveness of Vildagliptin in pediatric patients under 18 years of age have not been established.

**Drug Interactions** : As a result of studies no clinically relevant interactions with other oral antidiabetics (glibenclamide, pioglitazone, metformin), amlodipine, digoxin, ramipril, simvastatin, valsartan or warfarin were observed after co-administration with vildagliptin.

**Overdose** : Muscle pain, mild and transient paraesthesia, fever, oedema and transient increase in lipase levels.

**Storage** : Store below 30°C in a dry place.

**Packing : Dialiptin-50** : Each box contains 3 x 14's tablets in blister pack.