

Composition : Glucotin Max: Each Film coated tablet contains Glucosamine Sulfate 750mg as Glucosamine Sulfate Potassium Chloride USP and Diacerein INN 50mg.

Description : Glucosamine is a natural substance produced in the human body and found in certain foods. Glucosamine stimulates the production of glycosaminoglycan & proteoglycan, two essential building blocks of cartilage. Glucosamine acts, as a supplement, as a joint lubricant and shock absorber necessary to maintain healthy cartilage and joint function. Diacerein is an anti-inflammatory analgesic and antipyretic drug, indicated specially for the treatment of arthritic conditions. It directly inhibits IL-1 synthesis and release in vitro and down modulates IL-1 induced activities and relieves Osteoarthritis.

Indications : This tablet is indicated in the treatment of Osteoarthritis, Rheumatoid arthritis, Bone & Joint Injuries.

Dosage and administration : Adult and children 12 years and older : The usual dose of Glucosamine & Diacerein combination is 1 tablet twice daily (12 hourly) taken with food. Or, as directed by the registered physician.

Contraindication : It is contraindicated for the patient who have hypersensitivity to any component of the product.

Precautions : Caution should be taken in person who have diabetes, asthma, alcohol dependence or liver disease.

Side effects : Diarrhea, constipation, epigastric pain, heartburn, nausea, vomiting, headache, skin rashes, drowsiness, intense yellow coloring of urine may be associated with the active components of this product.

Glucotin Max

Tablet



Use in pregnancy and Lactation :

Glucosamine & Diacerein should not be administered during pregnancy. In addition, Diacerein should not be prescribed to lactating women due to reports that small amounts of Diacerein derivatives pass into the maternal milk.

Use in Child : This Medicine is not indicated for under 12 years of age.

Drug interactions : Decreased absorption of Diacerein with aluminium and/or magnesium hydroxide antacids. Increased risk of diarrhea with laxatives, antibiotics. It should be avoid co-administration with fibers and phytic acids. Interactions between glucosamine and other prescription or over-the counter medicines or herbal/health supplements have not been reported.

Overdose : Diarrhea may occur.

Storage : Store below 30°C in a dry place.

Packing : Glucotin Max : Each box contains 3 x 7's tablets in blister pack.