

Composition : Cefadroxil 500mg Capsule & 125mg/5ml Suspension.

Indications : Cefadroxil is indicated for the treatment of patients with urinary tract infection caused by Escherichia coli, Proteus mirabilis and Klebsiella species. Skin and skin structure infections caused by staphylococci, Pharyngitis and Tonsillitis caused by streptococcus pyogenes (Group A betahemolytic streptococci).

Dosage & administration : Administered orally and with food.

Adults : Urinary Tract Infections : The usual dosage is 1gm (2 capsules) or 2gm (4-capsules) per day in single or divided doses. Skin and Skin

Structure Infections : The usual dosage is 1gm (2 capsules) per day in single or divided doses. **Pharyngitis and Tonsillitis :** The usual dosage is 1gm (2 capsules) per day in single or divided doses. For urinary tract infections, the recommended daily dosage for children is 30mg/kg/day in divided doses every 12 hours.

Contraindication : Sefanid is contraindicated in patients with known allergy to the cephalosporin group of antibiotics.

Sefanid
Capsule



DRUG
INTERNATIONAL
LTD.

Precautions : Cefadroxil should be used with caution in the presence of markedly impaired renal function. In patients with known or suspected renal impairment, careful clinical observation is required.

Side effects : Cefadroxil is well tolerated and its adverse events generally mild and transient. Most commonly reported adverse events are dyspepsia, nausea, vomiting, hepatic dysfunction, abdominal pain, neutropenia etc.

Use in pregnancy and lactation : Cefadroxil has been assigned to pregnancy category B by the FDA.

Drug interaction : There are no significant data available.

Packing :

Sefanid Capsule : 3 x10's capsules in blister pack.

Sefanid Suspension : 100ml Suspension.